



KEY WORDS

- ✓ Salmonella serotypes
- ✓ Antimicrobial resistance
- ✓ Lactic acid bacteria
- ✓ Virulence factors

CONTACT

E-MAIL:

yldz.mrv20@gmail.com


THESIS SUPERVISOR

TELEPHONE:

(+90 224) 294 08 54

E-MAIL:

serpilkahya@uludag.edu.tr



DETERMINATION OF CARRIAGE RATE AND ANTIBIOTIC SENSITIVITY OF SALMONELLA IN CATS AND DOGS

Merve YILDIZ

0000-0003-2883-2338

BURSA ULUDAG UNIVERSITY
GRADUATE SCHOOL OF HEALTH SCEINCES
VETERINARY MICROBIOLOGY DEPARTMENT
PhD PROGRAM

GRADUATION DATE: 30.05.2024

SUPERVISOR

PROF. DR. SERPİL KAHYA DEMİRBİLEK 0000-0001-6138-7163 BURSA ULUDAG UNIVERSITY GRADUATE SCHOOL OF HEALTH SCIENCES VETERINARY MICROBIOLOGY DEPARTMENT BURSA – TÜRKİYE



THESIS ABSTRACT

The aim of the study was to determine the prevalence, risk factors, some virulence factors, serotypes and antimicrobial resistance profiles of *Salmonella* in domestic dogs and cats in Turkey, to evaluate the public health risk and to make a macroscopic comparison of LAB in *Salmonella*-positive and negative animals.

Salmonella prevalence was 5.73% (9/157) in dogs and 0.0% (0/191) in cats, of which 8 isolates were cultured by ISO method and 5 by FDA method. S. Enteritidis (n=4), S. Kentucky (n=3) and S. Virchow (n=2) serotypes were all positive for invA and stn virulence genes. MDR rate was 11.1% and the highest resistance was found for ciprofloxacin. MDR resistant S. Virchow and carbapenem resistant S. Enteritidis were detected.

APPLICATION AREAS OF THE THESIS RESULTS

There is limited data on Salmonella carriage in dogs and cats in Turkey. This study showed that apparently healthy dogs are subclinical carriers of Salmonella and harbor serotypes that cause human salmonellosis. In particular, aminoglycosides, monobactam and polymyxin antibiotics may be useful in the treatment of most cases of salmonellosis in dogs. Among zoonotic bacterial species such as Salmonella, isolation of carbapenem-resistant S. Enteritidis is a major public health problem. It is also important that all isolates harbor the virulence genes analyzed. The fact that Salmonella positive dogs consumed raw chicken meat emphasizes the association of Salmonella with raw meat consumption.

ACADEMIC ACTIVITIES

- Yıldız, M., & Kahya Demirbilek, S. (2023). Investigation of prevalence and antimicrobial resistance of Salmonella in pet dogs and cats in Turkey. Veterinary Medicine and Science, vms31513. https://doi.org/10.1002/vms3.1513
- Yıldız, M., & Demirbilek, S. K. (2023). Determination of Salmonella spp. prevalence and antibiotic resistance profiles in domestic animals. Journal of Istanbul Veterinary Sciences, 153-153. https://doi.org/10.2139/ssrn.4441458